

Rational Homotopy Theory

Joshua Moerman

Supervisor: Ieke Moerdijk


Introduction to homotopy theory

Rational homotopy theory

The main equivalence

Homotopy theory

Study of spaces or shapes
with “weak equivalences”



Important spaces

$$S^1 = \text{circle}$$

$$S^2 = \text{sphere}$$


$$S^3 = \dots$$

⋮


Important tool

Fundamental group:

$\pi_1(X) = \text{maps } S^1 \rightarrow X \text{ up to homotopy}$



$\pi_1(X) = \text{trivial}$



$\pi_1(Y) \neq \text{trivial}$

Important tools

Homotopy groups:

$\pi_1(X) = \text{maps } S^1 \rightarrow X \text{ up to homotopy}$

$\pi_2(X) = \text{maps } S^2 \rightarrow X \text{ up to homotopy}$

$\pi_3(X) = \text{maps } S^3 \rightarrow X \text{ up to homotopy}$

\vdots

Torsion-free

Serre proved in 1950s:

$$\text{odd } k : \quad \pi_n(S^k) \otimes \mathbb{Q} = \begin{cases} \mathbb{Q} & \text{if } n = k \\ 0 & \text{otherwise} \end{cases}$$

$$\text{even } k : \quad \pi_n(S^k) \otimes \mathbb{Q} = \begin{cases} \mathbb{Q} & \text{if } n = k, 2k - 1 \\ 0 & \text{otherwise} \end{cases}$$

Introduction to homotopy theory

Rational homotopy theory

The main equivalence

Rational homotopy theory

Study of spaces
with “rational equivalences”
and “rational homotopy groups”

Rational homotopy theory


Study of spaces
with “rational equivalences”
and “rational homotopy groups”

or

Study of *rational* spaces
with weak equivalences
and ordinary homotopy groups

Rational spaces

X is *rational* if $\pi_n(X)$ is a \mathbb{Q} -vector space



Introduction to homotopy theory

Rational homotopy theory

The main equivalence

Main equivalence

Theorem

Homotopy theory of rational spaces
=
*Homotopy theory of commutative differential
graded algebras*

Main equivalence (precise version)

Theorem

$$\mathbf{Ho}(\mathbf{Top}_{\mathbb{Q},1,f}) \simeq \mathbf{Ho}(\mathbf{CDGA}_{\mathbb{Q},1,f})^{op}$$

What is a cdga?

Definition

a cdga A is

- ▶ a \mathbb{Q} -vector space
- ▶ with a multiplication $A \otimes A \xrightarrow{\mu} A$
- ▶ with a differential $A \xrightarrow{d} A$ such that $d^2 = 0$
- ▶ with a grading $A = \bigoplus_{n \in \mathbb{N}} A^n$
- ▶ it is commutative: $xy = (-1)^{|x| \cdot |y|} yx$

Free cdga's

As always: there is a free guy: $\Lambda(\dots)$

For example

$$\Lambda(t, dt) \text{ with } |t| = 0$$

is just polynomials in t , with its differential dt

Dictionary

rational spaces

$S_{\mathbb{Q}}^n$ with n odd

$S_{\mathbb{Q}}^n$ with n even

Eilenberg-MacLane
space $K(\mathbb{Q}, n)$

cdga's

$\Lambda(e)$ with $|e| = n$

$\Lambda(e, f)$ with $|e| = n$,
 $|f| = 2n-1$ and $df = e^2$

$\Lambda(e)$ with $|e| = n$

Dictionary

rational spaces

weak equivalence

$$\pi_n(f) : \pi_n(X) \cong \pi_n(Y)$$

homotopy

$$h : X \times I \rightarrow Y$$

cdga's

weak equivalence

$$H(f) : H(X) \cong H(Y)$$

homotopy

$$h : A \rightarrow B \otimes \Lambda(t, dt)$$

Dictionary

rational spaces

$$\pi_n(X) = [S^n, X]$$

Long exact sequence of
a fibration

cdga's

$$\begin{aligned}\pi^n(A) &= H(Q(A)) \\ \pi^n(A)^* &\cong [A, \Lambda(e)] \\ &\text{or } [A, \Lambda(e, f)]\end{aligned}$$

Long exact sequence of
a cofibration

Dictionary


topological n -simplex

$$\Delta^n = \left\{ (x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid \sum x_i = 1, x_i \geq 0 \right\}$$


cdga n -simplex

$$\Delta_n = \frac{\Lambda(x_0, \dots, x_n, dx_0, \dots, dx_n)}{\langle \sum x_i - 1, \sum dx_i \rangle}, \quad |x_i| = 0$$

Construction



Construction



$$A(X) = \mathbf{Hom}_{\mathbf{sSet}}(X, \Delta(-))$$